

NOTAS SOBRE
MAMÍFEROS
SUDAMERICANOS

●



Sociedad Argentina para el Estudio de los Mamíferos



**Use of *Mauritia flexuosa* (Arecaceae) leaves as day roost by the
Dwarf Little Fruit bat *Rhinophylla pumilio* (Phyllostomidae)
in Mato Grosso, Brazil**

Guilherme S. T. Garbino (1), Vitor Q. Piacentini (2), Rogerio Vieira Rossi (2) y Thiago B. F. Semedo (3)

(1) Programa de Pós-Graduação em Zoologia, Laboratório de Mastozoologia, Instituto de Ciências Biológicas, Universidade Federal de Minas Gerais, Campus Pampulha, Belo Horizonte, MG, Brazil. (2) Departamento de Biologia e Zoologia, Instituto de Biociências, Universidade Federal de Mato Grosso, 78060-900, Cuiabá, MT, Brazil. (3) Programa de Capacitação Institucional, Museu Paraense Emílio Goeldi (MPEG) / Instituto Nacional de Pesquisa do Pantanal (INPP), Cuiabá, MT, Brazil. [correspondence: gstgarbino@hotmail.com]

ABSTRACT

We describe a group of Dwarf Little Fruit bats (*Rhinophylla pumilio*) using the Buriti palm (*Mauritia flexuosa*) as day roost in a Cerrado area in Cuiabá, Mato Grosso, Brazil. With ten individuals roosting under a dry Buriti palm leaf, we estimate that the group was composed of five females and their young. This is the first record of *R. pumilio* using leaves of *M. flexuosa*, and the third publication reporting the use of leaves of this palm as shelter by Phyllostomidae fruit bats.

RESUMO

Neste estudo, descrevemos um grupo de morcegos da espécie *Rhinophylla pumilio* utilizando a palmeira buriti (*Mauritia flexuosa*) como abrigo diurno em uma área de Cerrado em Cuiabá, Mato Grosso, Brasil. Com dez indivíduos empoleirados em uma folha seca de buriti, nós estimamos que este grupo era composto de cinco fêmeas e seus filhotes. Este é o primeiro registro do uso de folhas de *M. flexuosa* por *R. pumilio* e é a terceira publicação a reportar o uso de folhas de buriti como abrigo para morcegos Phyllostomidae.

The Dwarf Little Fruit bat, *Rhinophylla pumilio* Peters, 1865, is a small phyllostomid that lives in the humid forests of cis-Andean South America, from Colombia and Venezuela to southeastern Brazil. This frugivorous bat uses leaf tents and unmodified foliage as day roosts, but it may occasionally roost in culverts and inside buildings. In this note, we report the first case of *R. pumilio* using leaves of the Buriti palm (*Mauritia flexuosa*) as day roost in a Cerrado vegetation area. Based on the observed roosting group, we also comment on group behavior and reproduction of this bat species.

On December 11, 2019, while setting mist nets for birds, we found a group of *R. pumilio* roosting under a dry leaf of a *Mauritia flexuosa* (Arecaceae) palm. The leaf did not appear to have any modification, nor did it show signs of biting (Figure 1). The observation site is a riverine forest within a Cerrado vegetation type at Ecoville da Chapada, Cuiabá, Mato Grosso (Brazil), bordering Parque Nacional da Chapada dos Guimarães (15° 11' 21.0" S; 55° 56' 19.4" W; ca. 290 m above sea level).

Recibido el 22 de febrero de 2020. Aceptado el 4 de junio de 2020. Editor asociado: Mariano Sánchez.



A total of 10 bats were observed, but an individual flew away before the photograph was taken (Fig. 1). We identified the species as *R. pumilio* based on the central chin protuberance bordered by elongated fleshy pads, uniformly-colored noseleaf, brownish ears with whitish inner base, and olive-brown pelage. Our assumption that the photographed bats were *R. pumilio* is reinforced by the capture of an adult male on December 10, 2019, in the same area that the roosting group was recorded, which showed the same external characteristics described above. The collected specimen was captured using ground-level mist-nets, and handled following the guidelines of the American Society of Mammalogists (Sikes et al. 2016). It was deposited as a fluid-preserved specimen in the mammal collection of the Coleção Zoológica da Universidade Federal de Mato Grosso, Cuiabá, under the number UFMT 4890. The forearm length of the captured specimen is 34.2 mm, there is no diastema between I2 and C, the I1 has a cingular style, and the uropatagium is not conspicuously hairy, allowing us to determine it as *R. pumilio*. Additionally, *R. pumilio* is the only non-Stenodermatinae phyllostomid known to regularly use foliage as roosts.

Because parental care in *Rhinophylla pumilio* is carried out exclusively by the mother, we assumed that the larger animals with olive-brown pelage were females, and the smaller individuals with grayish pelage were their young. If the individual that flew away is accounted for, the roosting group consisted of ten bats, of which five were adult females and five were juveniles (we discard the possibility of the animal that escaped being a male because the group was left with five juveniles and four adults). The bats were huddled in a central position relative to the leaf lamina, and close to the leaf's petiole (Fig. 1).

Day roosts of *R. pumilio* in unmodified foliage are documented in at least 16 plant species, of which four are palm trees (Arecaceae): *Astrocaryum sciophilum*, *Attalea ataleoides*, *Jessenia bataua*, and *Mauritia flexuosa* (Table 1). We only found two other mentions of *Mauritia* leaves used as roosts by frugivorous leaf-nosed bats in the literature: an observation of *Artibeus lituratus* roosting under *Mauritia* sp. leaves in an urban area in southeastern Brazil, and records of unoccupied umbrella tents in *M. flexuosa* leaves on the island of Trinidad.

Considering adult individuals only, the size of the group recorded here is within the expected for *R. pumilio*. In French Guiana, group sizes ranged from 2 to 7 individuals, and in southeastern Brazil limited observations have recorded only roosting pairs.

Reproductive data for the Dwarf Little Fruit bat is scarce, and no clear pattern can be defined for the species. We assume that the females reported in this study were lactating due to the presence of younger animals attached to them. The lactating females of this study were recorded in the rainy season (early December), therefore births must have occurred by November/December.

In the Amazon of south-western Colombia, pregnant individuals have been recorded in May and July, and lactating individuals were captured in April, June, and December. In the Amazon of southern Venezuela, pregnant or lactating *R. pumilio* were found in December. Both in the Colombian and Venezuelan localities, there is a fairly constant rainfall volume throughout the year, so reproduction would not be correlated with the climatic season. In the Amazon of Brazil, pregnancies have been recorded in March and June, and lactating females in the dry season in August. In Espírito Santo, southeastern






Figure 1. Group of Dwarf Little Fruit bats (*Rhinophylla pumilio*) roosting under a Buriti palm (*Mauritia flexuosa*) dry leaf. A) lateral view of the leaf; B) ventral view of the leaf showing the group of *R. pumilio* with four adults (ad) and five juveniles (ju), and a detail of the face of one of the adult bats.

Brazil, pregnant females were captured during the rainy season between December and January. Our reproductive data suggests a pattern in central Brazil similar to what has been observed in southeastern Brazil, with births occurring in the rainy season when fruits are most abundant.

Most of what is known about the ecology of the Dwarf Little Fruit bat is from rain-forest areas. As fruit seasonality is an important factor in determining the reproductive patterns of tropical bats, more studies on the ecology of frugivorous bats from

Table 1. Foliage roosts used by *Rhinophylla pumilio* throughout its distribution, with the case reported herein marked in bold.

Plant species	Family	Roost type	Locality	Reference
<i>Philodendron fragrantissimum</i>	Araceae	leaf tent	French Guiana, Nouragues	Henry and Kalko (2007)
<i>P. melionii</i>	Araceae	leaf tent	French Guiana, Nouragues and St. Elie	Charles-Dominique (1993)
<i>P. ornatum</i>	Araceae	leaf tent	French Guiana, Nouragues	Henry and Kalko (2007)
<i>P. ornatum</i>	Araceae	leaf tent	French Guiana, St. Elie and Nouragues	Charles-Dominique (1993)
<i>Rhodospata latifolia</i>	Araceae	leaf tent	French Guiana, Nouragues	Charles-Dominique (1993)
<i>Attalea attaleoides</i>	Arecaceae	unmodified leaf	French Guiana, St. Elie	Charles-Dominique (1993)
<i>A. attaleoides</i>	Arecaceae	leaf tent	French Guiana, St. Elie	Charles-Dominique (1993)
<i>Atrocaryum sciophilum</i>	Arecaceae	leaf tent	French Guiana, Paracou, Nouragues, and St. Elie	Charles-Dominique (1993); Simmons and Voss (1998)
<i>A. sciophilum</i>	Arecaceae	unmodified leaf	French Guiana, Nouragues	Henry and Kalko (2007)
<i>Jessenia bataua</i>	Arecaceae	unmodified young leaf	French Guiana, Nouragues	Henry and Kalko (2007)
<i>Mauritia flexuosa</i>	Arecaceae	unmodified dry leaf	Brazil, Mato Grosso, Cuiabá	This study
Bromeliaceae sp. indet.*	Bromeliaceae	unmodified leaf	French Guiana, Nouragues	Charles-Dominique (1993)
Cyclanthaceae sp. indet.*	Cyclanthaceae	unmodified leaf	French Guiana, Nouragues	Charles-Dominique (1993)
<i>Heliconia</i> sp.	Heliconiaceae	leaf tent	Brazil, Espírito Santo	Zortéa (1995)
<i>Sterculia</i> sp.	Malvaceae	leaf tent	French Guiana, Nouragues	Charles-Dominique (1993)
<i>Musa</i> sp.	Musaceae	leaf tent	Brazil, Espírito Santo	Zortéa (1995)
<i>Phenakospermum guyannense</i>	Strelitziaceae	leaf tent	French Guiana, Paracou	Simmons and Voss (1998)
<i>P. guyannense</i>	Strelitziaceae	unmodified leaf	French Guiana, Paracou	Simmons and Voss (1998)
<i>Cecropia obtusa</i> *	Urticaceae	unmodified dry leaf	French Guiana, Nouragues	Charles-Dominique (1993)
<i>C. sciadophylla</i>	Urticaceae	unmodified dry leaf	French Guiana, Nouragues	Henry and Kalko (2007)

* used as roost after experimental removal of leaf tents (Charles-Dominique 1993).



the drier and more seasonal Cerrado habitats are necessary, to verify the existence of intraspecific differences between populations living in areas with different climates.

ACKNOWLEDGEMENTS

GSTG received a pre-doctoral fellowship from the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior, Brazil (CAPES; Code 001). TBFS is supported by a scholarship from the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) and the Ministério da Ciência, Tecnologia, Inovações e Comunicações (MCTIC) (process 300617/2020-8). We are thankful to the two anonymous reviewers and to the editor, for their constructive comments.

LITERATURE CITED

- GARBINO, G. S. T., & V. DA C. TAVARES. 2018. Roosting ecology of Stenodermatinae bats (Phyllostomidae): Evolution of foliage roosting and correlated phenotypes. *Mammal Review* 48: 75–89.
- HENRY, M., & E. K. V. KALKO. 2007. Foraging strategy and breeding constraints of *Rhinophylla pumilio* (Phyllostomidae) in the Amazon lowlands. *Journal of Mammalogy* 88: 81–93.
- KUNZ, T. H., & G. F. McCracken. 1996. Tents and harems: apparent defense of foliage roosts by tent-making bats. *Journal of Tropical Ecology* 12: 121–137.
- MARINKELLE, C. J., & A. CADENA. 1972. Notes on bats new to the fauna of Colombia. *Mammalia* 36: 50–58.
- McLELLAN, L. J., & K. F. KOOPMAN. 2008. Subfamily Carollinae Miller, 1924. *Mammals of South America, Volume 1: Marsupials, xenarthrans, shrews, and bats* (A. L. Gardner, ed.). The University of Chicago Press, Chicago, IL and UK.
- MORELLATO, L. P. C., M. G. G. CAMARGO, & E. GRESSLER. 2013. A review of plant phenology in South and Central America. *Phenology: An Integrative Environmental Science* (M. D. Schwartz, ed.). Springer, New York.
- PERACCHI, A. L., & S. T. ALBUQUERQUE. 1993. Quirópteros do município de Linhares, estado do Espírito Santo, Brasil (Mammalia, Chiroptera). *Revista Brasileira de Biologia* 53: 575–581.
- PERACCHI, A. L., S. D. L. RAIMUNDO, & A. M. TANNURE. 1984. Quirópteros do Território Federal do Amapá, Brasil (Mammalia, Chiroptera). *Arquivos da Universidade Federal Rural do Rio de Janeiro* 7: 89–100.
- REIS, N. R., & A. L. PERACCHI. 1987. Quirópteros da região de Manaus, Amazonas, Brasil (Mammalia, Chiroptera). *Boletim do Museu Paraense Emílio Goeldi* 3: 1–21.
- RINEHART, J. B., & T. H. KUNZ. 2006. *Rhinophylla pumilio*. *Mammalian Species* 791: 1–5.
- RODRÍGUEZ-HERRERA, B., R. A. MEDELLÍN, & R. M. TIMM. 2007. Murciélagos neotropicales que acampan en hojas. Editorial INBio, Santo Domingo.
- SAZIMA, I., W. A. FISCHER, M. SAZIMA, & E. A. FISCHER. 1994. The fruit bat *Artibeus lituratus* as a forest and city dweller. *Ciencia e Cultura* 46: 164–168.
- SIKES, R. S., & ASM ANIMAL CARE AND USE COMMITTEE. 2016. Guidelines of the American Society of Mammalogists for the use of wild mammals in research and education. *Journal of Mammalogy* 97: 663–688.
- SIMMONS, N. B., & R. S. VOSS. 1998. The Mammals of Paracou, French Guiana: a neotropical lowland rainforest fauna Part 1. Bats. *Bulletin of the American Museum of Natural History* 237: 1–219.
- SOLARI, S. ET AL. 2019. Family Phyllostomidae (New World Leaf-nosed Bats). *Handbook of the Mammals of the World, Bats, Vol. 9* (D. E. Wilson & R. A. Mittermeier, eds.). Lynx Edicions, Barcelona.
- TUTTLE, M. D., & D. STEVENSON. 1982. Growth and survival of bats. *Ecology of bats* (T. H. Kunz, ed.). Plenum Press, New York.
- WILSON, D. E. 1979. Reproductive patterns. *Biology of the bats of the New World Family Phyllostomatidae. Part III* (R. J. Baker, J. Knox Jones & D. C. Carter, eds.). Texas Tech Press, Lubbock, TX.
- ZORTÉA, M. 1995. Observations on tent-using in the Carolline bat *Rhinophylla pumilio* in Southeastern Brazil. *Chiroptera Neotropical* 1: 2–4.

