

NOTAS SOBRE MAMÍFEROS SUDAMERICANOS



Sociedad Argentina para el Estudio de los Mamíferos





Predation of *Leptodactylus macrosternum* (Anura: Leptodactylidae) by the Fringe-lipped bat, *Trachops cirrhosus* (Chiroptera: Phyllostomidae) in southeastern Brazil

Fred V. de Oliveira (1), Rodolfo Stumpp (2), and Maria Clara Nascimento-Costa (1)

(1) Laboratório de Evolução de Mamíferos, Instituto de Ciências Biológicas, Universidade Federal de Minas Gerais. Belo Horizonte, Minas Gerais, Brazil. (2) Laboratório de Ecologia e Conservação, Instituto de Ciências Biológicas, Universidade Federal de Minas Gerais. Belo Horizonte, Minas Gerais, Brazil. [correspondence: fredvictor97@gmail.com]

ABSTRACT

Although *Trachops cirrhosus* (Spix, 1823) is known for feeding on frogs, most of the data on its feeding behavior comes from the analysis of stomach contents, remains on roosting sites or experimental feeding in captivity, with few cases observed directly in the field. Here we report *T. cirrhosus* preying upon the frog *Leptodactylus macrosternum* Miranda-Ribeiro, 1926, which is the first record of this bat species preying an anuran in southeastern Brazil, and also the first direct observation of a predation event by *T. cirrhosus* in the wild in this region, contributing to the understanding of its trophic relations.

RESUMO

Embora o morcego *Trachops cirhrosus* seja conhecido por se alimentar de anuros, a maioria das informações sobre seus hábitos alimentares é resultado da análise de conteúdo estomacal, restos encontrados em abrigos ou alimentação experimental em cativeiro, com poucos casos observados diretamente na natureza. Neste trabalho nós reportamos *T. cirrhosus* predando a rã *Leptodactylus macrosternum*, sendo o primeiro registro deste morcego se alimentando de anuros no sudeste do Brasil, e também a primeira observação direta de um evento de predação por *T. cirrhosus* na natureza para esta região, contribuindo, desta forma, para o entendimento das suas relações tróficas.

Trachops cirrhosus (Spix, 1823), popularly known as Fringe-lipped bat, is a medium-sized Phyllostomidae occurring in forest and stream habitats, open savannas and agricultural areas from southern Mexico to southern Brazil (Cramer et al. 2001, Williams & Genoways 2007, Miller et al. 2015). This species is closely related to animalivorous and omnivorous species in the family Phyllostomidae (Baker et al. 2016), and it is easily distinguishable from other bat species due to conical or cylindrical wart-like protuberances on its chin and lips, serrated edge on the noseleaf, and a large pinna (Cramer et al. 2001).

The diet of *T. cirrhosus* has been reported to include insects, fruits and small vertebrates such as birds, lizards, and even other bats (e.g., Peracchi et al. 1982; Bonato & Facure 2000; Rodrigues et al. 2004; Leal et al. 2018). However, the species is mainly known from its ability to catch and feed on frogs, and being capable of using prey-generated acoustic cues to discriminate among poisonous and palatable Recibido el 29 de febrero de 2020. Aceptado el 13 de mayo de 2020. Editor asociado: Agustín Abba.

prey (Ryan & Tuttle 1983). Although T. cirrhosus includes a wide range of items in its diet, few data are available on its feeding habits and the knowledge of its ecology is still poorly known, especially in Brazil (Rodrigues et al. 2004; Leal et al. 2018). Within this scenario, the goals of this study were to report a new predation event of T. cirrhosus upon the anuran Leptodactylus macrosternum Miranda-Ribeiro, 1926 in southeastern Brazil, contributing to the understanding of its trophic relations and natural history.

Fieldwork was conducted in July 2017 (dry season) at Fazenda Fortaleza de Santa Terezinha, a livestock farm in the municipality of Jequitaí, state of Minas Gerais, southeastern Brazil (17° 10' 44.78" S; 44° 38' 41.80"W). The native vegetation cover in the area is mainly composed of dry forests ("Mata Seca"), and narrow gallery and riparian forests on the banks of rivers and streams within the Cerrado biome (sensu IBGE 2004).

On July 11, 2017, we installed six ground-level mist-nets (each net is 12 m long and 3 m high) on a trail within a small fragment of gallery forest along a dry watercourse. One male specimen of T. cirrhosus (forearm 61.09 mm, head-body 73 mm, weight 40.13 g) was captured close to the ground (approximately at a height of 1 m), while carrying an already dead and half-eaten frog, identified as L. macrosternum (Anura: Leptodactylidae) (Fig. 1A, B). Several individuals of the same frog species were found in the nearby forest where the mist net was placed (Fig. 1C). The individual T. cirrhosus was collected under the license IEF 034.012/2016/MG, and all capture and collection procedures followed the CFBio (Conselho Federal de Biologia) and IBAMA (Instituto Brasileiro do Meio Ambiente e dos Recursos Naturais Renováveis) resolutions. The specimen was deposited at the Coleção de Mamíferos do Centro de Coleções Taxonômicas da Universidade Federal de Minas Gerais (CM-CCT/UFMG) under the catalog number UFMG 6811 and had tissue samples collected.

Most data on the feeding behavior of T. cirrhosus comes from stomach content analyses, remains on roosting sites or experimental feeding in captivity, with few cases observed in the field (Cramer et al. 2001; Leal et al. 2018). This is the first direct observation of a predation event by T. cirrhosus in the wild in southeastern Brazil and also the first record of this bat species preying an anuran in this region. Other records on the diet of the Fringe-lipped bat in Brazil are restricted to Amazonia (northern Brazil), and the states of Pernambuco (northeastern Brazil) and Rio de Janeiro (southeastern Brazil) (Peracchi et al. 1982; Rodrigues et al. 2004; Rocha et al. 2012; Rocha et al. 2016; Castro & Costa-Campos 2017; Leal et al. 2018). Nonetheless, data from Rio de Janeiro were obtained from captive bats and include items that may not be usual for the species in the wild (e.g., Mus musculus). In Brazil, amphibians were reported in the diet of T. cirrhosus by Bonato et al. (2004), Rocha et al. (2012), Rocha et al. (2016), Castro & Costa-Campos (2017), and Leal et al. (2018). Particularly, frogs of the genus Leptodactylus were reported on the ground of feeding perches of T. cirrhosus in northeastern Brazil (Leal et al. 2018) and this is the second record of this genus in the diet of Trachops. The consumption we report herein occurred during the dry season when the consumption of amphibians by T. cirrhosus would be less expected since these bats use frogs' call to locate its prey, and frog vocal activity peak occur mostly on the rainy season (Duellman & Trueb 1994; Zina & Haddad 2005). Although anurans are cited as common prey of the Fringe-lipped bat, Leal et al. (2018) showed insects are the main item in the diet of *T. cirrhosus*, followed by amphibians. Silva (2007) also reports insects as the most frequent items on fecal samples of *T. cirrhosus*, followed by fruits and small vertebrates. These findings show that, although *T. cirrhosus* may be able to feed on frogs and small vertebrates, it is a generalist predator as other phyllostomines.

ACKNOWLEDGMENTS

We would like to thank D. Santana and H. C. Costa for helping with the frog's identification, H. C. Costa for the review and comments on the original manuscript and R. Costa for both the photographs of the *Leptodactylus*.



Figure 1. Specimen of *Trachops cirrhosus* (UFMG 6811) captured at Fazenda Fortaleza de Santa Terezinha, Jequitaí, Minas Gerais state, Brazil (A), predating on Leptodactylus macrosternum (B). Live specimen of *Leptodactylus macrosternum* commonly found in the area (C). Scale bar=1cm

LITERATURE CITED

- BAKER, R. J., S. SOLARI, A. CIRRANELO, & N. B. SIMMONS. 2016. Higher level classification of Phyllostomid bats with a summary of DNA synapomorphies. Acta Chiropterologica 18: 1–38.
- BONATO, V., & K. G. FACURE. 2000. Bat predation by the fringe-lipped bat *Trachops cirrhosus* (Phyllostomidae, Chiroptera). Mammalia 64: 241–243.
- BONATO, V., K. G. FACURE, & W. UIEDA. 2004. Food habits of bats of subfamily Vampyrinae in Brazil. Journal of Mammalogy 85: 708–713.
- CASTRO, I. J., & C. E. COSTA-CAMPOS. 2017. New record of anuran predation by *Trachops cirrhosus* (Mammalia, Chiroptera, Phyllostomidae) in a varzea forest in the estuary of the Amazon River, Eastern Amazon. Biota Amazônia 7: 108–109.
- CRAMER, M. J., M. R. WILLIG, & C. JONES. 2001. Trachops cirrhosus. Mammalian Species 656: 1-6.
- DUELLMAN, W. E., & TRUEB, L. 1994. Biology of Amphibians. Johns Hopkins University Press, Baltimore/ London.
- LEAL, E. S. B. ET AL. 2018. What constitutes the menu of *Trachops cirrhosus* (Chiroptera)? A review of the species' diet. Neotropical Biology and Conservation 13: 337–346.
- MILLER, B., F. REID, J. ARROYO-CABRALES, A. D. CUARÓN, & P. C. DE GRAMMONT. 2015. Trachops cirrhosus. The IUCN Red List of Threatened Species 2015: e.T22029A22042903. http://dx.doi.org/10.2305/ IUCN.UK.2015-4.RLTS.T22029A22042903.en
- PERACCHI, A. L., S. T. E ALBUQUERQUE, & S. D. L. RAIMUNDO. 1982. Contribuição ao conhecimento dos hábitos alimentares de *Trachops cirrhosus* (Spix, 1823) (Mammalia, Chiroptera, Phyllostomidae). Arquivos da Universidade Federal Rural do Rio de Janeiro 5: 1–5.
- ROCHA, R., I. SILVA, A. M. DOSREIS, & G. M. ROSA. 2012. Another frog on the menu: predation of *Trachops cirrhosus* (Chiroptera: Phyllostomidae) upon *Osteocephalus oophagus* (Anura: Hylidae). Chiroptera Neotropical 18: 1136–1138.
- ROCHA, R., M. GORDO, & A. LÓPEZ-BAUCELLS. 2016. Completing the menu: Addition of Scinax cruentommus and Scinax cf. garbei (Anura: Hylidae) to the diet of Trachops cirrhosus (Chiroptera: Phyllostomidae) in Central Amazon. North-Western Journal of Zoology 12: 199–204.
- RODRIGUES, F. H. G., M. L. REIS, & V. S. BRAZ. 2004. Food habits of the Frog-eating Bat, *Trachops cirrhosus*, in Atlantic Forest of northeastern Brazil. Chiroptera Neotropical 10: 180–182.
- RYAN, M. J., & M. D. TUTTLE. 1983. The ability of the Frog-eating bat to discriminate among novel and potentially poisonous frog species using acoustic cues. Animal Behaviour 31: 827–833.
- SILVA, L. A. M. 2007. Comunidades de morcegos em uma área de Caatinga e Brejo de Altitude no agreste de Pernambuco. Brasília, DF. Tese de Doutorado. Universidade de Brasília, Brazil.
- WILLIAMS, S. L. & H. H. GENOWAYS. 2007. Subfamily Phyllostominae Gray, 1825. Mammals of South America Volume I Marsupials, Xenarthrans, Shrews, and Bats (A. L. Gardner, ed.). The University of Chicago Press, Chicago and London.
- ZINA, J., & C. F. B HADDAD. 2005. Reproductive activity and vocalizations of *Leptodactylus labyrinthicus* (Anura: Leptodactylidae) in Southeastern Brazil. Biota Neotropica 5: 1–11

